

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE October 1999		3. REPORT TYPE AND DATES COVERED September 1999	
4. TITLE AND SUBTITLE ACCEPTABILITY OF CARBOHYDRATE DRINKS DURING THE MARINE CORPS INFANTRY OFFICER COURSE 10-DAY FIELD TRAINING EXERCISE				5. FUNDING NUMBERS	
6. AUTHOR(S) Scott J. Montain, Susan M. McGraw, William J. Tharion, F. Matthew Kramer, and Reed W. Hoyt					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Nutrition and Biochemistry Division and Thermal and Mountain Medicine Division U.S. Army Research Institute of Environmental Medicine Bldg. 42 Natick, Massachusetts 01760-5007				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Same as block #7				10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES					
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release - distribution unlimited				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Soldiers typically under eat relative to their caloric expenditure during field training. Carbohydrate supplements in liquid and solid form have been developed to provide a means of enhancing carbohydrate and total caloric intake when soldiers are likely to consume insufficient rations. This report summarizes the results of a taste test performed during a seven-day field training exercise. Marines were provided five packets of ERGO Drink (Soldier Systems Center-Natick, Soldier Biological Chemical Command, Natick, Massachusetts) per day during the training exercise. They completed a questionnaire at the end of the 7-day sampling period. The soldiers rated the ERGO Drink as very good and thought it was a very valuable product for use in the field, particularly when relying primarily on MRE for food. The packaging was also highly rated. The students chose to dissolve the powder either in their canteen or within the packet itself. On other occasions they consumed the drink as a powder. The canteen cup was generally not utilized for fluid delivery. These results document the usefulness and acceptability of carbohydrate beverage powder for field use. Novel means of packaging should be pursued to encourage consumption directly from the drink packet, thus avoiding contamination of the canteen with carbohydrate.					
14. SUBJECT TERMS fluid replacement, exercise, diet, field rations, carbohydrate, sports drinks, sustained operations				15. NUMBER OF PAGES 37	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT		

19991202 123

USARIEM TECHNICAL REPORT

ACCEPTABILITY OF CARBOHYDRATE DRINK SUPPLEMENTATION DURING THE MARINE CORPS INFANTRY OFFICER COURSE 10-DAY FIELD TRAINING EXERCISE

Scott J. Montain, Ph.D.

Susan M. McGraw, B.S.

William J. Tharion, M.S.

F. Matthew Kramer, Ph.D.

Reed W. Hoyt, Ph.D.

Military Nutrition and Biochemistry Division

Thermal and Mountain Medicine Division

and

Supporting Science and Technology Directorate, Soldier Systems Center - Natick

October 1999

U.S. Army Research Institute of Environmental Medicine

Natick, MA 01760-5007

TABLE OF CONTENTS

LIST OF FIGURES	iii
ACKNOWLEDGEMENTS	iv
EXECUTIVE SUMMARY	1
INTRODUCTION	1
METHODS	2
Subjects	2
Experimental Design	3
Statistical Analysis	4
RESULTS	4
Subject Characteristics	4
Acceptability of ERGO Drink	7
Acceptability of ERGO Drink Packaging	10
Methods of Consumption	12
Utility of Carbohydrate Liquid Supplements for Use in Military Activities	15
Overall Rating of ERGO Drink	15
Written Comments	16
DISCUSSION	17
CONCLUSIONS	22
RECOMMENDATIONS	22
REFERENCES	24
APPENDIX A. CARBOHYDRATE DRINK QUESTIONNAIRE.....	26

LIST OF FIGURES

Figure 1. Nutritional supplement use	6
Figure 2. Mean (filled star) and median (unfilled x) responses regarding ERGO Drink acceptability	7
Figure 3. Frequency distribution of selected ERGO Drink acceptability parameters.....	8
Figure 4. Acceptability of ERGO Drink flavors, value as a ration supplement, and likelihood of consumption if available	9
Figure 5. Acceptability of ERGO Drink packaging	11
Figure 6. Methods of ERGO Drink consumption during 7 days of field training	13
Figure 7. Behavioral strategies for consuming the ERGO Drink during the 7-day field exercise.....	14
Figure 8. Expected use during military training and overall like or dislike.....	16

ACKNOWLEDGEMENTS

The authors wish to thank the Marines of the Infantry Officer Course, Class 5-99, of The Basic School, Marine Corps Combat Development Command, Quantico, Virginia, for their participation in the study; the Marine Corps Basic School for their wonderful and dedicated support prior to and during data collection; The US Marine Corps for their financial support; SGTs' Kevin Warren and Mark Buller for their technical support; and Ms. Theresa Skibinski and Ms. Shelley Strowman for their help in developing the carbohydrate drink questionnaire.

EXECUTIVE SUMMARY

Soldiers typically under eat relative to their caloric expenditure during field operations. Carbohydrate supplements in liquid and solid form have been developed to provide a means of enhancing carbohydrate and total caloric intake when soldiers are likely to consume insufficient rations. This report summarizes the results of an acceptability test performed during a seven-day field exercise. Seventy-nine Marines were provided five packets of ERGO Drink (Soldier Systems Center, US Army Soldier Biological Chemical Command, Natick, Massachusetts) per day during the operation. They completed a questionnaire at the end of the operation. The Marines rated the ERGO Drink as very good and thought it was a very valuable product for use in the field, particularly when relying primarily on Meals, Ready-to-Eat (MRE) as the primary source of food. The packaging was also highly rated. The Marines were divided in their preferred method of consumption, choosing to either dissolve the powder in their canteen or within the packet itself, and occasionally consuming the drink as a powder. The canteen cup was not utilized as means of consumption. These results document the acceptability and advisability of carbohydrate beverage powder for field use. Novel means of packaging should be pursued to encourage consumption directly from the drink packet, thus avoiding contamination of the canteen with carbohydrate.

INTRODUCTION

Soldiers working in field environments typically under eat relative to their caloric expenditure (Baker-Fulco, 1995). This behavior results in not only inadequate caloric intake but also inadequate carbohydrate intake to maintain carbohydrate energy reserves and physical performance (Montain et al., 1997). Recently, scientists at the Soldier Systems Center, US Army Soldier Biological and Chemical Command, Natick, Massachusetts developed a carbohydrate beverage base designed to supplement field rations. This item, termed the "ERGO Drink" is a 12% mass/volume blend of maltodextrin, glucose and fructose (9%, 2%, and 1%, respectively). Five flavors are available (Lemon, Lemon-lime, Raspberry, Tropical Fruit, and Punch). It is packed in a 9x13 cm pouch containing 47 g of carbohydrate (170 kilocalories). When diluted to proper strength, the package provides a 0.35 liter (12 ounce) beverage. This report summarizes the acceptability and utility of this carbohydrate supplement for field use.

METHODS

SUBJECTS

Seventy-nine male students in Infantry Officer Course, Class 5-99, The Basic School, Marine Corps Combat Development Command, Quantico, Virginia 22554, volunteered to participate in this study. During the sampling period, they were participating in a 10-day field exercise. The Marines were informed of the purpose and methodology of the experiment prior to beginning the field exercise.

EXPERIMENTAL DESIGN

During the first seven days of the ten-day field exercise, the Marines were provided a sufficient quantity of ERGO Drink to enable each Marine the opportunity to consume 5 ERGO Drink packets per day. They were issued the packets prior to the field exercise and during one re-supply point during the course. No restrictions were placed on the number of drinks per day, time of day of consumption, or method of consumption. The Marines were simply asked to consume the drinks during the exercise and provide feedback via a questionnaire after seven days of sampling.

The field exercise consisted of daily force-on-force operations, designed to develop leadership skills during combat-like conditions. Sleep periods ranged from several hours per day during the initial days of the operations to short naps during the final days of the exercise. Daily activities included planning and preparation, combat maneuvers, and de-briefing. The daily caloric expenditure in an earlier class was approximately 6,000 kilocalories per day. The students were provided five Meals, Ready-to-Eat (MRE) at the beginning of the field exercise and at one re-supply point, thus limiting caloric intake from MRE to approximately 1300 kilocalories per day.

ERGO Drink flavors provided were Lemon and Tropical Fruit. No verbal instructions were given regarding how to prepare the beverage base, but the package instructs the user to "mix the contents of one package in 12 fluid ounces of water. Stir until the powder dissolves."

Following completion of the course, Marines completed a questionnaire (Appendix B) regarding acceptance, utility and preparation issues.

STATISTICAL ANALYSIS

Means, medians, and frequency distributions were calculated using SPSS statistical software. Chi squared analysis was performed to identify flavor preference. Data are presented as mean \pm sd. When mean and median responses are presented together, the following format is used: mean \pm sd; median.

RESULTS

SUBJECT CHARACTERISTICS

The Marines averaged 25 \pm 2 years of age (range 22-33 yr.). Their self-reported supplement use is presented in Figure 1. Forty-eight of the 79 (61%) Marines reported that they were nutritional supplement users. Twenty-eight of the Marines (35%) reported taking vitamin and/or mineral supplements at least once per week. Fewer Marines (13 of 79; 16%) reported taking amino acid or protein supplements (including creatine) once or more per week.

The Marines were familiar with sports drinks, sports bars and carbohydrate gels. Twenty-six of 79 (33%) reported drinking carbohydrate sports drinks one or more times per week. An additional ten Marines consumed them on an occasional basis. Thirty-nine of 79 Marines (49%) ate sports bars at least occasionally, while 24 of 79 Marines (30%) had consumed carbohydrate gels prior to the course.

The Marines had mixed opinions on the adequacy of the field rations as only 37 of 79 (47%) felt that their nutritional needs were met by the MRE. The most frequent written comments were that the field rations did not contain enough food and/or calories (n=27), and that the ration was either too high in fat or did not contain enough complex

carbohydrates, protein, and vegetables (n = 16). It is noteworthy that the Marines unanimously thought that supplements should be added to the current ration system.




Figure 1. Nutritional supplement use.

ACCEPTABILITY OF ERGO DRINK

Figure 2 presents the mean and median responses of the group regarding product acceptability. The appearance, flavor, smell, sweetness, and overall acceptance were quite high. Figures 3 and 4 present the frequency distributions of the same data. The Marines only slightly to moderately liked the appearance and smell of the ERGO Drink; largely due to many Marines providing a neutral response. Flavor and sweetness scored higher marks with both mean and median scores ranging from "like moderately" to "like very much." Lemon Flavor and Tropical Fruit Flavor scored even higher marks with both mean and median scores ranging from "like moderately" to "like very much."




Figure 2. Mean (filled star) and median (unfilled x) responses regarding ERGO Drink acceptability.




Figure 3. Frequency distribution of selected ERGO Drink acceptability parameters.

Both flavors received favorable scores. Seventy percent of the volunteers rated the lemon flavor ERGO Drink a 7 to 9 on the 9-point scale (moderately like to extremely like). Ninety percent of the Marines rated the tropical fruit flavor a 7 to 9 on the same scale. Chi square analysis revealed no difference in drink flavor acceptability ($\chi^2 = 1.57$).

The majority of Marines thought the ERGO Drink would be a valuable ration supplement, with 85% of responses within the “very likely” to “extremely likely” categories. When queried about the likelihood of consuming the ERGO Drink if it was available as a supplement to their rations, 73% selected “extremely likely” and none picked a response below “neutral.”




Figure 4. Acceptability of ERGO Drink flavors, value as a ration supplement, and likelihood of consumption if available.

ACCEPTABILITY OF ERGO DRINK PACKAGING

Figure 5 illustrates the frequency distribution of responses regarding the ERGO Drink packaging. The package size appeared appropriate for field operations, as it was rated "just right" in size (2.75 ± 0.79 ; 3.0 on 5 point scale). The package was rated as "very easy" to open (7.9 ± 1.1 ; 8.0) as well as "moderately" to "very easy" to use (7.5 ± 1.4 ; 8.0). The Marines generally felt the portion size was adequate, but would prefer a somewhat larger size (6.0 ± 1.1 ; 6.0). Seventy-eight of 79 Marines (99%) preferred a soft package to a hard package. Sixty-three of 79 (80%) of the Marines preferred powder form to dissolvable paste ($n = 6$; 8%) or tablet ($n = 10$; 13%).







Figure 5. Acceptability of ERGO Drink packaging.

METHODS OF CONSUMPTION

To assess the method of ERGO Drink consumption, the Marines were asked a single multi-part question in which they were to rate the proportion of time they either consumed the ERGO Drink as a powder (straight from package), dissolved in their canteen cup, dissolved in their canteen, diluted in the ERGO Drink package, or by some other means. Fifty-nine of 79 Marines (75%) completed the questionnaire correctly. The remaining 20 questionnaires had certain options unmarked, didn't properly account for time (e.g., "one method always" and another "some of the time") or the question was left unanswered. Of these 20 questionnaires, 9 were dropped from the analysis and the other 11 were modified to properly account for time. If a soldier stated that they always used one method and sometimes another (the most frequent incorrect response), the "always" was modified to "most of the time." Similarly, if the Marines had neglected to mark "never" on an option, but relative proportion of use for other options was adequate to account for 100% of use, "never" was marked for that option. Figure 6 summarizes the methods of ERGO Drink consumption for the 59 appropriately filled out answers and the 11 modified questionnaires.




Figure 6. Frequency distribution of the individual methods of ERGO Drink consumption during 7 days of field training.

It is clear that very few Marines chose to dilute the ERGO Drink in their canteen cup. Instead, the methods of choice were either to dissolve the drink powder in their canteen, add water directly to the ERGO Drink pouch, or consume the product as a dry powder. Twenty-nine of 68 respondents (43%) chose to dissolve the ERGO Drink in their canteen at least $\frac{1}{2}$ time. Thirty-one of 68 respondents (46%) chose to add water to the ERGO Drink packet $\frac{1}{2}$ time or more. The other method used at least "some of the time" was to eat the powder straight from the package. This latter method was used by

51 of 67 (76%) Marines at least occasionally, and 25% used it as their primary means of intake.

Most Marines chose more than one method to consume the ERGO Drink. Figure 7 illustrates the relative usage for those individuals who reported dissolving the ERGO Drink in their canteen, dissolving it in the ERGO package, or eating it as a dry powder. Most common (24%), was to chose all three methods of consumption. Next most common were to consume the ERGO Drink as a dry powder, at least part of the time and to either dissolve it in the canteen (19%) or dissolve the powder inside the ERGO Drink package (17%) the rest of the time.




Figure 7. Behavioral strategies for consuming the ERGO Drink during the 7-day field exercise.

UTILITY OF CARBOHYDRATE LIQUID SUPPLEMENTS FOR USE IN MILITARY ACTIVITIES

The frequency distributions regarding expected use of carbohydrate drinks in selected military venues are presented in Figure 8. The expected use in garrison was varied among the Marines, with the most frequent response being "somewhat likely". Expected use in field venues was higher, reaching a mean numeric score of 7.1 ± 1.8 and 8.7 ± 0.7 when relying on field kitchen and MREs, respectively. These numeric scores equate to "moderately likely" and "very likely". Marines, furthermore, expected that the drink would be most commonly consumed during physical training (mean = 8.0 ± 1.4) vs. during rest (mean = 7.0 ± 1.9) or with meals (mean = 7.1 ± 1.9).

OVERALL RATING OF ERGO DRINK

The ERGO Drink received an overall score of 8.1 ± 0.9 which equates to "like very much". Frequency distribution of individual responses is shown in Figure 8. Sixty-six of 79 (84%) rated the ERGO Drink "like very much" or "like extremely". No one surveyed rated the ERGO Drink unfavorably.




Figure 8. Expected use during different training environments and overall like or dislike.

WRITTEN COMMENTS

The Marines had the opportunity to provide written comments after several of the questions. Twenty-two of those surveyed stated that they thought the ERGO Drink should be added to the MRE and/or will use the product again if it becomes available. When queried about additional/alternate flavors, 10 suggested grape, 5 suggested cherry, and 4 each recommended orange and strawberry. As for improving the product,

the most frequent recommendation ($n = 10$) was to increase package size so that the beverage can be prepared in its package. Less frequent ($n = 4$) was a complaint that the 2-ply package comes apart or is flimsy. Other suggestions received more than once recommended: easier dissolution, a bar and/or gel form, packet quantity appropriate for 1 Qt. canteen, resealable package and/or easier to consume on the move.

DISCUSSION

The results of this evaluation during repeated days of field exercises demonstrate the high acceptability of the ERGO Drink as a ration supplement. The two ERGO Drink flavors examined, tropical fruit and lemon, were both given very favorable hedonic ratings, and the product was given an overall score of 8 on a 9-point scale.

The results also demonstrate that military personnel will use and appreciate this type of product during field training exercises. The Marines verbally told the scientific staff that they looked forward to drinking the ERGO Drink each day and when queried about the likelihood of using this product if it were available, the ERGO Drink received a remarkable 9 on the 9-point scale.

The ERGO Drink's flavor, smell and sweetness also scored very well. These high marks occurred despite the fact that for most of the time, the drink was consumed in a form more concentrated than recommended. This could mean that the Marines in this study may have preferred sweet tasting items. Alternatively, the high proportion of maltodextrin in the beverage may have limited excessive sweetness when it was consumed in more concentrated form. The use of maltodextrin provides the ERGO Drink a possible advantage over other products relying on glucose, sucrose and

fructose for carbohydrate, since the ERGO Drink features significantly more carbohydrate without excessive sweetness. Whether Marines would rate other products equal to, lower or higher than the ERGO Drink remains untested as is the question of whether there is any physical and/or cognitive advantage of consuming the ERGO Drink over commercial sports drinks.

The multi-day assessment has clear advantages over a single sampling sessions as it enabled evaluation of the ERGO Drink over many hours of work and many repeated samplings. Repeated use also provided the opportunity for the Marines to determine the "best" method of field consumption – an important issue since the product is expected to be used most by physically active troops with limited access to food or time to eat. It is noteworthy that the ERGO Drink was so favorably received as it might have been expected that the two flavors would become monotonous over an extended sampling time. Whether the distinctness of the two flavors enhanced the acceptability of each individually and/or reduced potential monotony is unknown. When considering the extremely high hedonic ratings, it must be acknowledged that rations were purposefully restricted during the course, despite high levels of physical activity. Caloric restriction may therefore have inflated the ratings of the ERGO Drink compared to assessment under more calorically balanced situations. Nevertheless, one MRE per day for dismounted forces, who commonly carry in excess of 45 kg, is a realistic daily food intake for operational forces.

Whether the ERGO Drink improved the cognitive and physical capabilities of the Marines was not examined. However, several course instructors commented that the class appeared less physically and mentally strained compared to previous classes.

Furthermore, unlike previous classes, the Marines in this study never developed the qualitative signs of ketosis.

Despite the high acceptability scores given the ERGO Drink, the results are restricted to the two flavors tested. While we would have preferred to examine the acceptability of all five flavors, the other three flavors were not available for evaluation at the time of the study. Future studies should evaluate the acceptability of the remaining flavors.

The ERGO Drink packaging also received above average marks. Important to highly mobile troops, the package was rated as very easy to open and use. While portion size was acceptable, there was some interest in making the portion size somewhat larger. This could have, in part, been driven by the interest of some individuals for packaging the beverage for a 1-Qt container rather than the ½ Qt canteen cup.

The two most common methods of consumption were as a slurry directly from the ERGO Drink package and as dry powder. Seventy percent of the Marines chose to mix the powder in the ERGO Drink package at least some of the time and 42% chose this method "most of the time" or "always". This suggests that modifications to the ERGO Drink package should be considered to make the package more usable as a drink container. Several Marines provided written comments suggesting that the package be made somewhat larger and less flimsy. The majority (99%) of Marines preferred it remain a soft package.

Few of the soldier chose to mix the ERGO Drink in their canteen cup. Fifty-seven of 65 Marines who answered the question (87%) never mixed the ERGO Drink in

the canteen cup and only one soldier used this method more than ½ of the time. Cup usage was likely limited due to the inconvenience of removing the canteen cup from the canteen holder, reluctance to clean the canteen cup, and the availability of more convenient means to ingest the product. Due to the unpopularity of this method, some modification to the instructions should be considered, as it is unlikely that the product will be used in the canteen cup.

It is important to note that 68% (47 of 69) of the Marines used their canteen to dissolve the ERGO Drink and 33% (23 of 69) used this as their primary method of ingesting the drink. This raises concerns about potential water container contamination, as well as lack of water for wound cleaning. It is well recognized that the addition of carbohydrate will promote mold and bacterial growth, while some flavorings can impair water purification technologies. Efforts should be taken to reduce the likelihood of carbohydrate drinks being stored in the canteen (e.g., education, a warning label on the packet) or making canteen sanitation a high priority for military personnel working in field environments

Additional improvements to the beverage powder and/or packaging should also be considered. As many Marines mixed the product inside the ERGO Drink packet, water was added to the product. In such conditions, maltodextrin does not readily enter into solution but rather remains lumpy. One solution would be to alter the packaging so that the powder could be added to water in a separate packet.

The drink would also have greater utility for field use if it contained electrolytes consistent with the concentrations recommended by the National Academy of Sciences (Committee on Military Nutrition Research, 1994). While the ERGO Drink was

developed as a carbohydrate supplement rather than a fluid-electrolyte replacement product, there is little reason that it could not serve both purposes. The addition of sodium, chloride, and potassium, in quantities closely approximating those amounts lost in sweat (15-30 mEq/L sodium, 15-30 mEq/L chloride, 2-5 mEq/L potassium) would provide an electrolyte source when ration intake may be insufficient to replace daily electrolyte losses. During work periods with a high water turnover and little electrolyte replacement, soldiers and athletes are under the greatest risk for heat cramps, salt-depletion heat exhaustion, and clinical symptoms associated with hyponatremia (Ladell, 1957; Garigan and Ristedt, 1999; Speedy et al., 1999). The latter condition, which is potentially life threatening, can be avoided during prolonged work by ingestion of small amounts of salt. If the ERGO Drink contained electrolytes, it could serve as a source of both carbohydrates and electrolytes for persons likely to under-eat during field operations. Additionally, small amounts of sodium chloride would enable the drink to serve as an oral rehydration solution during periods of gastrointestinal illness, a condition common to military operations.

The ERGO Drink was developed as a liquid carbohydrate supplement to combat the under-feeding which occurs during military field training. As a beverage, however, it can also be a helpful fluid replacement solution as persons will voluntarily drink more of a flavored, sweetened drink than water alone (Hubbard et al., 1984). Under most military situations, a 12% solution is acceptable as both a carbohydrate energy source and as a fluid replacement source. Because of the ERGO Drink's relatively high concentration of carbohydrate, it slightly compromises gastric emptying and rehydration during periods of profuse sweating (for review, Coyle and Montain, 1991). The

relatively lower sweating rates elicited by military scenarios versus sports competition, makes this a smaller concern. Furthermore, during situations in which high rates of fluid replacement are required to sustain performance, the ERGO Drink can be diluted in half to optimize carbohydrate and water absorption.

CONCLUSIONS

The ERGO Drink was given high approval ratings by troops working in a field. The sweetness, flavoring, packaging and ease of use combined to recommend it as an extremely valuable supplement to field rations.

RECOMMENDATIONS

1. Further refinements in product packaging should be pursued, such as a package that can be also used as a cup, as the majority of Marines chose to dilute the product directly in the package or consume it dry rather than use their canteen cup.
2. Electrolytes should be added to the ERGO Drink to make it a more versatile beverage for hot weather field use.
3. The remaining ERGO Drink flavors should be examined for acceptability during field training.
4. Additional studies should be performed to:
 - a. Examine interaction of multiple flavors on acceptability and use during field operations.
 - b. Examine acceptability and use when rations are not purposefully restricted.
 - c. Examine acceptability of different forms of packaging during field training.

- d. Examine the influence of ERGO Drink availability on mission performance during sustained operations.
 - e. Examine the influence of ERGO Drink on nutrient intake during field training.
5. The ERGO Drink should be compared to commercial sports beverages for acceptability and ergogenic effects.

REFERENCES

Baker-Fulco, C.J. An overview of dietary intakes during military exercises. In: Not Eating Enough, B.M. Marriot (Ed) National Academy Press, Washington, DC: 121-146, 1995.

Committee on Military Nutrition Research. Committee Summary and Recommendations. In: Fluid replacement and heat stress, B.M. Marriot (Ed) National Academy Press, Washington, DC: 3-8, 1994.

Coyle, E.F., Montain, S.J. Carbohydrate and fluid ingestion during exercise: are there tradeoffs? Med Sci Sports Exerc 24: 671-678, 1992.

Garigan, T., Ristedt, D.E. Death from hyponatremia as a result of acute water intoxication in an Army Basic Trainee. Mil Med 164: 234-237, 1999.

Hubbard, R.W., Sandick, B.L., Matthew, W.T., Francesconi, R.P., Sampson, J.B., Durkot, M.J., Maller, O., Engell, D.B. Voluntary dehydration and alliesthesia for water. J Appl Physiol: 57: 868-875, 1984.

Ladell, W.S.S. Disorders due to heat. Trans Roy Soc Trop Med Hyg 51: 189-216, 1957.

Montain, S.J., Shippee, R.L., Tharion, W.J. Carbohydrate – electrolyte solution effects on physical performance of military tasks. Aviat Space Environ Med 68: 384-391, 1997.

Speedy, D.B., Noakes, T.D., Rogers, I.R., Thompson, J.M.D, Campbell, R.G.D., Kuttner, J.A., Boswell, D.R., Wright, S., Hamlin, M. Hyponatremia in ultradistance athletes. Med Sci Sports Exerc 31: 809-815, 1999.

APPENDIX A. CARBOHYDRATE DRINK QUESTIONNAIRE

Carbohydrate Beverage Acceptability Survey

MARKING INSTRUCTIONS

- Use a No. 2 pencil only.
- Do not use ink, ballpoint, or felt tip pens.
- Make solid marks that fill the response completely.
- Erase cleanly any marks you wish to change.
- Make no stray marks on this form.

CORRECT: ●

INCORRECT: ☐ ⊗ ⊖ ⊙

ID NUMBER			
0	0	0	0
1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

DATE			
MONTH	DAY	YEAR	
<input type="radio"/> Jan			
<input type="radio"/> Feb			
<input type="radio"/> Mar	0	0	0
<input type="radio"/> Apr	1	1	1
<input type="radio"/> May	2	2	2
<input type="radio"/> June	3	3	3
<input type="radio"/> July	4	4	4
<input type="radio"/> Aug	5	5	5
<input type="radio"/> Sept	6	6	6
<input type="radio"/> Oct	7	7	7
<input type="radio"/> Nov	8	8	8
<input type="radio"/> Dec	9	9	9

Product Acceptability

1. Using the scale below, please rate, overall, how much you liked or disliked the following aspects of the carbohydrate beverage.

	Did Not Eat/Drink	Dislike Extremely	Dislike Very Much	Dislike Moderately	Dislike Slightly	Neither like nor Dislike	Like Slightly	Like Moderately	Like Very Much	Like Extremely
Appearance	0	1	2	3	4	5	6	7	8	9
Flavor	0	1	2	3	4	5	6	7	8	9
Smell	0	1	2	3	4	5	6	7	8	9
Sweetness	0	1	2	3	4	5	6	7	8	9

Comment(s): _____

Do not write
in this box

0	1	2	3	4	5	6	7	8	9
0	1	2	3	4	5	6	7	8	9

2. Please rate how much you liked or disliked the following flavor(s). Please fill in a bubble for each flavor.

	Did Not Eat/Drink	Dislike Extremely	Dislike Very Much	Dislike Moderately	Dislike Slightly	Neither like nor Dislike	Like Slightly	Like Moderately	Like Very Much	Like Extremely
Lemon/Lime	0	1	2	3	4	5	6	7	8	9
Lemon	0	1	2	3	4	5	6	7	8	9
Tropical Fruit	0	1	2	3	4	5	6	7	8	9
Punch	0	1	2	3	4	5	6	7	8	9
Orange	0	1	2	3	4	5	6	7	8	9
Raspberry	0	1	2	3	4	5	6	7	8	9

What other flavors would you like? _____

Do not write
in this box

0	1	2	3	4	5	6	7	8	9
0	1	2	3	4	5	6	7	8	9

Product Packaging**3. Using the scale below, please rate the package size.**

Much Too Small Somewhat Too Small Just Right Somewhat Too Large Much Too Large

1	2	3	4	5
---	---	---	---	---

4. Using the scale below, please rate the ease of opening the package.

Extremely Difficult Very Difficult Moderately Difficult Somewhat Difficult Neither Easy or Difficult Somewhat Easy Moderately Easy Very Easy Extremely Easy

1	2	3	4	5	6	7	8	9
---	---	---	---	---	---	---	---	---

5. Please rate the ease of using the product.

Extremely Difficult Very Difficult Moderately Difficult Somewhat Difficult Neither Easy or Difficult Somewhat Easy Moderately Easy Very Easy Extremely Easy

1	2	3	4	5	6	7	8	9
---	---	---	---	---	---	---	---	---

6. Would you prefer the portion size to be:

Much Smaller Quite A Bit Smaller Moderately Smaller Somewhat Smaller Neither Smaller or Larger Somewhat Larger Moderately Larger Quite A Bit Larger Extremely Larger

1	2	3	4	5	6	7	8	9
---	---	---	---	---	---	---	---	---

7. What type of packaging would you prefer for this product? Please fill only one response.

- ☐ Soft Package
☐ Hard Package

8. In what form would you prefer this product? Please fill only one response.

- ☐ Powder
☐ Dissolvable Tablet
☐ Dissolvable Paste

9. Would the carbohydrate beverage be a valuable supplement to your field rations?

Extremely Unlikely Very Unlikely Moderately Unlikely Somewhat Unlikely Neither likely or unlikely Somewhat Likely Moderately Likely Very Likely Extremely Likely

1	2	3	4	5	6	7	8	9
---	---	---	---	---	---	---	---	---

Product Utility

10. Would you eat/drink this item if it were available as a supplement to your field rations?

Extremely Unlikely Very Unlikely Moderately Unlikely Somewhat Unlikely Neither likely or unlikely Somewhat Likely Moderately Likely Very Likely Extremely Likely

1	2	3	4	5	6	7	8	9
---	---	---	---	---	---	---	---	---

11. Using the scale below, please rate how likely you would be to use this product during:

Extremely Unlikely Very Unlikely Moderately Unlikely Somewhat Unlikely Neither likely or unlikely Somewhat Likely Moderately Likely Very Likely Extremely Likely

Garrison training	1	2	3	4	5	6	7	8	9
-------------------	---	---	---	---	---	---	---	---	---

Field training:

When field kitchen is available	1	2	3	4	5	6	7	8	9
---------------------------------	---	---	---	---	---	---	---	---	---

When MRE is primary food source	1	2	3	4	5	6	7	8	9
---------------------------------	---	---	---	---	---	---	---	---	---

12. Using the scale below, please rate how likely you would be to use the carbohydrate beverage during:

Extremely Unlikely Very Unlikely Moderately Unlikely Somewhat Unlikely Neither likely or unlikely Somewhat Likely Moderately Likely Very Likely Extremely Likely

Physical Activity	1	2	3	4	5	6	7	8	9
-------------------	---	---	---	---	---	---	---	---	---

Rest Periods	1	2	3	4	5	6	7	8	9
--------------	---	---	---	---	---	---	---	---	---

Meal Time	1	2	3	4	5	6	7	8	9
-----------	---	---	---	---	---	---	---	---	---

13. Overall, how much do you like or dislike the carbohydrate beverage?

Did Not Eat Dislike Extremely Dislike Very Much Dislike Moderately Dislike Slightly Neither like nor Dislike Like Slightly Like Moderately Like Very Much Like Extremely

0	1	2	3	4	5	6	7	8	9
---	---	---	---	---	---	---	---	---	---

14. What is your age today?

Please write your response in the blank boxes, then fill in the corresponding circles.

age		1	2	3	4	5	6				
		0	1	2	3	4	5	6	7	8	9

15. What is your gender?

M
F

Male

Female

16. Are you a nutritional supplement user?

Y
N

Yes

No (please go to question number 17) Never/hardly Once in a while 1-6 times per week Once a day

Vitamins/minerals	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Amino acids/protein (including creatine)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Herbal Preparations (e.g., ginseng, garlic, etc.)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Carbohydrate-Electrolyte Beverages	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Carbohydrate Sports Bars	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Carbohydrate Gels	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Other _____	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Do not write
in this box

0 1 2 3 4 5 6 7 8 9
0 1 2 3 4 5 6 7 8 9

17. Do the current field rations meet your nutritional needs?

Y
N

Yes (please go to question number 18)

No

In NO, why not? _____

Do not write
in this box

0 1 2 3 4 5 6 7 8 9
0 1 2 3 4 5 6 7 8 9

18. Do you think that nutritional supplements should be added to field rations?

Y
N

Yes

No

19. Do you think the carbohydrate beverage could improve your performance?

Y
N

Yes

No

Do you have any suggestions for improving the product? _____

Do not write
in this box

0 1 2 3 4 5 6 7 8 9
0 1 2 3 4 5 6 7 8 9

20. During this field exercise, did you consume the ERGO drink as:

	Never	Some of Time	Half of Time	Most of Time	Always
Dry Powder	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Dissolved in Canteen Cup	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Dissolved in Canteen	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Dissolved in ERGO package	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Other	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

If 'other', please describe: _____

21. Additional Comments: _____

Thank You!

DISTRIBUTION LIST

1 Copy:

Defense Technical Information Center
8725 John J. Kingman Road, Suite 0944
Fort Belvoir VA 22060-6218

Office of the Assistant Secretary of Defense (Health Affairs)
ATTN: Medical Readiness, Pentagon
Washington DC 20310-0103

Deputy Director for Medical Readiness
The Joint Staff, J-4
ATTN: J4-MRD
4000 Joint Staff, Pentagon
Washington DC 20318-4000

HQDA, Assistant Secretary of the Army
(Research, Development and Acquisition)
ATTN: SARD-TM, Pentagon
Washington DC 20316-0103

HQDA, Office of The Surgeon General
ATTN: DASG-RDZ
Room 3E368, Pentagon
Washington DC 20310-2300

HQDA, Office of The Surgeon General
ATTN: DASG-ZA
5109 Leesburg Pike
Falls Church VA 22041-3258

HQDA, Office of The Surgeon General
ATTN: DASG-DB
5109 Leesburg Pike
Falls Church VA 22041-3258

HQDA, Office of The Surgeon General
ATTN: DASG-MS
5109 Leesburg Pike
Falls Church VA 22041-3258

HQDA, Office of The Surgeon General
ATTN: SGPS-PSP, Preventive Medicine Consultant
5109 Leesburg Pike
Falls Church VA 22041-3258

U.S. Army Medical Research and Materiel Command
ATTN: MCMR-OP
504 Scott Street
Fort Detrick MD 21702-5012

U.S. Army Medical Research and Materiel Command
ATTN: MCMR-PLC
504 Scott Street
Fort Detrick MD 21702-5012

U.S. Army Medical Research and Materiel Command
ATTN: MCMR-PLE
504 Scott Street
Fort Detrick MD 21702-5012

Uniformed Services University of the Health Sciences
ATTN: Dean, School of Medicine
4301 Jones Bridge Road
Bethesda MD 20814-4799

Uniformed Services University of the Health Sciences
ATTN: Chair, Department of Preventive Medicine
4301 Jones Bridge Road
Bethesda MD 20814-4799

Commandant
Army Medical Department Center and School
ATTN: HSMC-FR, Bldg. 2840
Fort Sam Houston TX 78236-6100

Commandant
Army Medical Department Center and School
ATTN: Chief Librarian, Stimson Library
Bldg. 2840, Room 106
Fort Sam Houston TX 78236-6100

Commandant
Army Medical Department Center and School
ATTN: Director of Combat Development
Fort Sam Houston TX 78236-6100

Commander
U.S. Army Aeromedical Research Laboratory
ATTN: MCMR-UAX-SI
Fort Rucker AL 36362-5292

Commander
U.S. Army Medical Research Institute
of Chemical Defense
ATTN: MCMR-UVZ
Aberdeen Proving Ground MD 21010-5425

Commander
U.S. Army Medical Materiel Development Activity
ATTN: MCMR-UMZ
504 Scott Street
Fort Detrick MD 21702-5009

Commander
U.S. Army Institute of Surgical Research
ATTN: MCMR-USZ
3400 Rayley E. Chambers Avenue
Fort Sam Houston TX 78234-5012

Commander
U.S. Army Medical Research Institute
of Infectious Diseases
ATTN: MCMR-UIZ-A
504 Scott Street
Fort Detrick MD 21702-5011

Director
Walter Reed Army Institute of Research
ATTN: MCMR-UWZ-C (Director for Research Management)
Washington DC 20307-5100

Commander
U.S. Army Soldier Systems Commander
ATTN: AMSSC-CG
Natick MA 01760-5000

Commander
U.S. Army Natick Research, Development
and Engineering Center
ATTN: SSCNC-Z
Natick MA 01760-5000

Commander
U.S. Army Natick Research, Development
and Engineering Center
ATTN: SSCNC-S-IMI
Natick MA 01760-5040

Commander
U.S. Army Natick Research, Development
and Engineering Center
ATTN: SSCNC-TM (U.S Marine Corps Representative)
Natick MA 01760-5004

Director
U.S. Army Research Institute for Behavioral Sciences
5001 Eisenhower Avenue
Alexandria VA 22333-5600

Commander
U.S. Army Training and Doctrine Command
ATTN: ATMD (Office of the Surgeon)
Fort Monroe VA 23651-5000

Commander
U.S. Army Center for Health Promotion
and Preventive Medicine
Aberdeen Proving Ground MD 21010-5422

Director, Biological Sciences Division
Office of Naval Research - Code 141
800 N. Quincy Street
Arlington VA 22217

Commanding Officer
Naval Medical Research & Development Command
NNMC/Bldg 1
Bethesda MD 20889-5044

Commanding Officer
U.S. Navy Clothing & Textile Research Facility
ATTN: NCTRF-01, Bldg 86
Natick MA 01760-5053

Commanding Officer
Naval Environmental Health Center
2510 Walmer Avenue
Norfolk VA 23513-2617

Commanding Officer
Naval Aerospace Medical Institute (Code 32)
Naval Air Station
Pensacola FL 32508-5600

Commanding Officer
Naval Health Research Center
Bethesda MD 20889

Commanding Officer
Naval Health Research Center
P.O. Box 85122
San Diego CA 92138-9174

Commander
Armstrong Medical Research Laboratory
Wright-Patterson Air Force Base OH 45433

U.S. Air Force Aeromedical Library
Document Services Section
2511 Kennedy Circle
Brooks Air Force Base TX 78235-5122

Commander
U.S. Air Force School of Aerospace Medicine
Brooks Air Force Base TX 78235-5000

Director
Human Research and Engineering Directorate
U.S. Army Research Laboratory
Aberdeen Proving Ground MD 21005-5001

U.S. Army Biomedical R&D Representative
Science and Technology Center, Far East
ATTN: AMC-S&T, FE. Unit 45015
APO 96343-5015

National Defence Headquarters
ATTN: Research and Development Branch
Human Performance Directorate
305 Rideau Street
Ottawa Ontario CANADA K1A 0K2

Department of National Defence
Defence Research Establishment Ottawa
ATTN: Head, Physiology Group
Environmental Sciences Section
Protective Sciences Division
Ottawa Ontario CANADA M3M 3B9

Director
Defence and Civil Institute of Environmental Medicine
1133 Sheppard Avenue W.
P.O. Box 200
Downsview Ontario CANADA M3M 3B9

Defence and Civil Institute of Environmental Medicine
ATTN: Head, Environmental Physiology Section
1133 Sheppard Avenue W.
P.O. Box 200
Downsview Ontario CANADA M3M 3B9

Back Cover